Birat Journal Of Health Sciences
ISSN (Online): 2542-2804 • Quarterly • Vol. 8, No. 2 (2022)

Association Between Maternal Oral Health and Adverse Pregnancy Outcomes

Timothy Dobbins[1]
[1] School of Population Health, Faculty of Medicine and Health, UNSW Sydney, Australia
Email: t.dobbins@unsw.edu.au

ARTICLE HISTORY

RECEIVED	REVISED	ACCEPTED	PUBLISHED ONLINE
[14 May 2022]	*[17 July 2022]*	*[22 August 2022]*	*[31 August 2022]*

Abstract

Maternal oral health represents an important component of comprehensive prenatal healthcare because periodontal infections, gingival inflammation, dental caries, and inadequate oral hygiene may influence maternal well-being during pregnancy. Increasing attention has been directed toward the potential association between maternal periodontal disease and adverse pregnancy outcomes, including preterm birth, low birth weight, preeclampsia, and fetal growth restriction. This narrative review examines the relationship between maternal oral health and pregnancy outcomes, emphasizing periodontal disease, inflammatory mechanisms, maternal risk factors, clinical evidence, preventive strategies, and interdisciplinary healthcare approaches. Evidence from observational studies and systematic reviews suggests that periodontal disease is associated with increased risks of certain adverse maternal and neonatal outcomes, although findings vary substantially across populations and study designs. Proposed mechanisms include systemic inflammatory responses, changes in maternal immune regulation, and possible dissemination of oral microorganisms or their products. However, associations do not establish causation, and randomized clinical trials have not consistently demonstrated that periodontal treatment during pregnancy prevents preterm delivery or other adverse obstetric outcomes. Maternal dental assessment, appropriate periodontal treatment, oral hygiene education, and collaboration between dental and obstetric professionals remain important for maintaining oral health during pregnancy. The review emphasizes that integrating oral healthcare into prenatal services may improve maternal dental health and access to treatment, while further research is required to clarify its potential influence on pregnancy outcomes.

Keywords: Maternal Oral Health, Periodontal Disease, Adverse Pregnancy Outcomes, Preterm Birth, Low Birth Weight, Preeclampsia, Gingival Inflammation, Prenatal Dental Care, Maternal Health, Oral Health Education

COPYRIGHT & LICENSE

Copyright © 2026 The Author(s). All rights reserved by the author(s), who retain copyright ownership of their original work. The author(s) grant the **Birat Journal of Health Sciences (BJHS)** a non-exclusive right of first publication.

Open Access License: This article is published under the [Creative Commons Attribution 4.0 International (CC BY 4.0) License](https://creativecommons.org/licenses/by/4.0/), permitting unrestricted use, distribution, reproduction, and adaptation in any medium, including commercial use, provided that appropriate credit is given to the original author(s), the source and the license, and any modifications are clearly indicated.

1. Introduction

Pregnancy is associated with significant physiological, hormonal, immunological, and metabolic changes that may influence oral health. Increased concentrations of reproductive hormones can alter gingival vascular responses and inflammatory reactions, potentially increasing susceptibility to gingival swelling and bleeding. Changes in dietary habits, nausea, vomiting, and oral hygiene practices

may further affect dental health during pregnancy. Consequently, maternal oral conditions deserve consideration as part of comprehensive prenatal healthcare. Periodontal disease is a chronic inflammatory condition affecting the supporting structures of the teeth. Its development is associated with microbial dental plaque and the host immune response. During pregnancy, periodontal inflammation may become more clinically apparent, particularly among women with preexisting gingival disease or inadequate plaque control. Chronic inflammation has attracted considerable research interest because of its possible relationship with systemic maternal conditions and pregnancy complications. Adverse pregnancy outcomes, including preterm birth, low birth weight, fetal growth restriction, and preeclampsia, contribute substantially to maternal and neonatal morbidity. These complications arise through multiple pathways involving maternal medical conditions, infections, placental function, environmental exposures, and socioeconomic factors. Researchers have investigated whether periodontal infection and oral inflammatory processes contribute additional risk or serve as markers of broader maternal health inequalities. Observational investigations have reported associations between periodontal disease and certain adverse pregnancy outcomes. However, findings are inconsistent, and some well-designed studies have not identified statistically significant relationships. Clinical trials evaluating periodontal treatment have also produced mixed results, emphasizing the importance of distinguishing epidemiological association from a demonstrated causal effect. Maternal oral healthcare remains important regardless of whether periodontal treatment prevents obstetric complications. Untreated dental disease can cause pain, infection, difficulties eating, and reduced quality of life. Improving oral health knowledge, providing access to appropriate dental services, and strengthening communication between dental and prenatal healthcare professionals may support maternal well-being. This article evaluates the association between maternal oral health and adverse pregnancy outcomes through six interconnected areas: pregnancy-related oral changes, biological mechanisms, evidence concerning preterm birth and low birth weight, maternal complications, preventive dental healthcare, and future research and policy directions.

Pregnancy-Related Physiological Changes and Maternal Oral Health Conditions

Pregnancy produces physiological changes that can influence the health of gingival tissues and the oral environment. Increased levels of estrogen and progesterone affect vascular permeability, tissue responses, and inflammatory processes. These changes may intensify gingival reactions to dental plaque, resulting in swelling, redness, tenderness, and bleeding. Pregnancy-associated gingivitis is therefore an important oral healthcare concern, although pregnancy itself does not inevitably produce periodontal disease. Existing plaque accumulation and inadequate oral hygiene remain important contributors to gingival inflammation (Silk et al., 2008). Periodontal disease involves inflammation affecting the gingiva and, in more advanced cases, destruction of the supporting tissues surrounding teeth. Gingivitis is generally reversible with appropriate plaque control, whereas periodontitis involves loss of periodontal attachment and may contribute to progressive deterioration of tooth-supporting structures. The distinction is clinically important because pregnancy-related gingival inflammation should not automatically be classified as destructive periodontal disease. Accurate assessment requires examination of gingival condition, periodontal attachment, probing depths, and other relevant clinical indicators (Sanz & Kornman, 2013). Maternal oral health can also be affected by changes in dietary preferences and eating frequency. Some pregnant women experience increased snacking or cravings for carbohydrate-rich foods, potentially increasing exposure to fermentable sugars. Nausea and vomiting may complicate oral hygiene practices, while repeated exposure to gastric acid can contribute to dental erosion. These factors may increase the risk of dental discomfort and oral disease when appropriate preventive care is unavailable. Oral healthcare counseling should therefore address dietary habits, toothbrushing practices, fluoride exposure, and management of nausea-related oral symptoms (Silk et al., 2008). Dental caries represents another important maternal oral health concern. Untreated cavities may lead to pain, pulpal infection, and difficulties maintaining adequate dietary intake. Although dental caries and periodontal disease have different biological mechanisms, they may coexist in individuals experiencing barriers to oral healthcare. Maternal oral

conditions can also influence the oral health environment of infants through bacterial transmission and shared family practices. Comprehensive prenatal dental assessment should therefore evaluate both periodontal and dental health rather than focusing exclusively on gum inflammation (Boggess & Edelstein, 2006). Access to dental treatment during pregnancy may be limited by misconceptions concerning the safety of professional oral healthcare. Some pregnant women postpone dental appointments because they believe that routine treatment may adversely affect fetal development. Healthcare professionals may also hesitate to recommend dental procedures without clear knowledge of pregnancy-related treatment considerations. Such beliefs can contribute to avoidable delays in addressing oral infections and dental discomfort. Evidence-based prenatal counseling should explain that appropriately indicated preventive and therapeutic dental care can generally be provided safely during pregnancy, with necessary clinical precautions (Silk et al., 2008). Maternal oral health assessment should include discussion of symptoms, previous dental treatment, oral hygiene habits, tobacco exposure, dietary factors, and existing medical conditions. Identifying periodontal inflammation early can facilitate appropriate preventive measures and dental referral. Women with significant gingival bleeding, periodontal attachment loss, dental infections, or persistent pain may require professional evaluation. Integrating these assessments into prenatal healthcare can promote maternal comfort and support continuity between dental and obstetric services (Boggess & Edelstein, 2006). Overall, pregnancy-related changes may increase the visibility of preexisting oral health problems and create additional challenges for maintaining dental hygiene. However, the severity of maternal oral disease is influenced by multiple factors, including plaque control, individual susceptibility, healthcare access, and socioeconomic circumstances. Recognition of these interacting determinants is essential when evaluating the potential relationship between maternal oral health and pregnancy outcomes (Sanz & Kornman, 2013).

Biological Mechanisms Linking Periodontal Inflammation With Pregnancy Complications

The potential association between maternal periodontal disease and adverse pregnancy outcomes has attracted scientific attention because chronic oral infection may contribute to systemic inflammatory activity. Periodontitis involves a complex interaction between pathogenic microorganisms and the host immune response. Inflammatory processes within periodontal tissues can produce mediators that enter the circulation and potentially influence distant physiological systems. Researchers have proposed that these processes may affect maternal vascular responses, placental function, and pregnancy-related inflammatory pathways (Offenbacher et al., 1996). Periodontal infections commonly involve microbial communities containing Gram-negative anaerobic bacteria and other microorganisms associated with chronic tissue inflammation. These organisms and their components can stimulate the production of inflammatory mediators, including interleukins, tumor necrosis factor, and prostaglandins. Such mediators are involved in normal immune responses but may contribute to pathological processes when inflammatory activity is excessive or persistent. Because inflammatory signaling also plays a role in cervical maturation and labor initiation, researchers have investigated whether chronic periodontal inflammation could influence the timing of delivery (Sanz & Kornman, 2013). One proposed mechanism involves the dissemination of bacterial products or inflammatory mediators from periodontal tissues into maternal circulation. Inflamed gingival tissues may provide opportunities for transient bacteremia during ordinary oral activities such as chewing or toothbrushing. Although transient bacteremia does not necessarily cause systemic disease, investigators have considered whether sustained periodontal inflammation might influence maternal immune responses. These hypotheses provide biological plausibility for an association between periodontal disease and obstetric complications, but direct causal pathways remain incompletely established (Xiong et al., 2007). Placental inflammation is another potential mechanism through which maternal oral infection has been investigated. The placenta plays a central role in fetal nutrition, oxygen exchange, and immunological communication between mother and fetus. Excessive inflammatory responses or vascular disturbances may impair placental function and contribute to complications such as fetal growth restriction or hypertensive disorders. Periodontal disease has been studied as a potential contributor to these processes, although available evidence cannot consistently

determine whether oral inflammation independently initiates placental abnormalities (Boggess et al., 2003). The proposed relationship between periodontal disease and preeclampsia has received particular attention. Preeclampsia is a complex pregnancy disorder characterized by new-onset hypertension after 20 weeks of gestation, accompanied by proteinuria or other evidence of maternal organ dysfunction. Endothelial dysfunction, abnormal placental development, and inflammatory processes are involved in its pathophysiology. Researchers have hypothesized that chronic periodontal inflammation could contribute to systemic inflammatory burden and vascular dysfunction. However, similarities between the inflammatory characteristics of the two conditions do not establish a direct causal relationship (Cota et al., 2006). An additional explanation involves shared risk factors rather than direct biological causation. Maternal smoking, nutritional status, diabetes, socioeconomic disadvantage, healthcare access, and other characteristics may influence both periodontal health and pregnancy outcomes. When such factors are distributed unevenly between study groups, they can create or exaggerate statistical associations. Even studies that adjust for known confounders may not fully account for differences in health behaviors, chronic disease severity, or healthcare utilization. Therefore, biological mechanisms must be interpreted alongside epidemiological limitations (Xiong et al., 2006). The relationship between oral inflammation and pregnancy may also be bidirectional or influenced by common physiological pathways. Pregnancy-related hormonal and immune changes can alter periodontal responses, while maternal systemic conditions may affect both oral tissue health and obstetric risk. Consequently, a statistical association between periodontal disease and an adverse outcome may reflect several interacting processes. Understanding these relationships requires longitudinal studies that establish temporal sequences and carefully evaluate alternative explanations (Sanz & Kornman, 2013). Biological research supports the plausibility of interactions between oral infections, systemic inflammation, and maternal physiological processes. Nevertheless, the strength of the evidence differs across proposed mechanisms. Future investigations should examine whether specific microbial profiles, inflammatory biomarkers, or periodontal disease characteristics predict pregnancy complications independently of established maternal risk factors (Xiong et al., 2007).

Association of Maternal Periodontal Disease With Preterm Birth and Low Birth Weight

Preterm birth and low birth weight are among the most frequently investigated pregnancy outcomes in relation to maternal periodontal disease. Preterm birth is generally defined as delivery before 37 completed weeks of gestation, whereas low birth weight refers to a birth weight below 2,500 grams. These outcomes are associated with increased risks of neonatal complications, although their underlying causes differ. Preterm birth concerns gestational timing, while low birth weight may result from prematurity, restricted fetal growth, or both. Research assessing maternal periodontal disease should therefore distinguish these outcomes rather than automatically combining them into a single measure (Xiong et al., 2006). An early case-control investigation examined the association between periodontal infection and preterm low birth weight among 124 women. The researchers reported substantially greater periodontal disease among mothers experiencing adverse birth outcomes than among controls. After adjustment for selected obstetric and behavioral factors, maternal periodontal disease remained statistically associated with preterm low birth weight. These findings stimulated considerable interest in oral infection as a potentially modifiable maternal health factor. However, the study's observational design and relatively small sample limited conclusions concerning causality (Offenbacher et al., 1996). Subsequent prospective research provided additional evidence supporting an association. A study involving 1,313 pregnant women evaluated periodontal health during the second trimester and examined subsequent birth outcomes. Participants with severe or generalized periodontitis demonstrated increased odds of preterm delivery after adjustment for selected maternal characteristics. The association was particularly pronounced for earlier preterm deliveries. Nevertheless, the investigators emphasized that observational findings could not determine whether periodontal disease directly caused premature birth (Jeffcoat et al., 2001). Further evidence emerged from research examining periodontal disease and preterm low birth weight among pregnant women. An investigation conducted in Chile reported a higher frequency of adverse birth outcomes among women with periodontal disease than among comparison groups. These findings contributed to the

hypothesis that maternal periodontal health could influence pregnancy outcomes. However, differences in treatment exposure, baseline characteristics, and study design require careful consideration when interpreting associations between oral disease and neonatal outcomes (López et al., 2002). Systematic reviews have attempted to determine whether the associations reported in individual studies remain consistent across populations. A review of observational evidence found that many investigations identified increased risks of preterm birth, low birth weight, or related complications among women with periodontal disease. Nevertheless, other studies reported no significant association, and differences in periodontal assessment methods limited direct comparison. The authors concluded that evidence suggested a possible relationship but was insufficient to establish a definitive causal connection (Xiong et al., 2006). Contrasting evidence has emerged from multicenter prospective investigations. A study involving 786 pregnant women, including 311 with periodontal disease and 475 without, examined a composite outcome involving preterm birth, preeclampsia, fetal growth restriction, or perinatal death. After adjustment for relevant confounding factors, the researchers found no statistically significant association between periodontal disease and the composite outcome or individual outcomes of preterm birth and preeclampsia. These results demonstrate that associations observed in some populations are not consistently reproduced across different study settings (Srinivas et al., 2009). Differences in periodontal disease definitions may contribute to inconsistent findings. Some studies classify disease according to probing depth, while others emphasize clinical attachment loss, bleeding, or combinations of periodontal indicators. The timing of oral examination may also influence observed associations because periodontal conditions can change during pregnancy. Studies that use inconsistent definitions may therefore classify women differently even when their clinical characteristics are similar. Standardized periodontal assessment is necessary to improve research comparability (Sanz & Kornman, 2013). The available evidence suggests that maternal periodontal disease may be associated with preterm birth and low birth weight in certain populations, but the magnitude and independence of these associations remain uncertain. Observational findings support continued investigation, while conflicting results emphasize the need to control for maternal risk factors and avoid interpreting correlation as proof of causation. Maternal oral healthcare should be promoted for established dental health benefits without promising prevention of premature birth (Xiong et al., 2007).

Evidence Concerning Maternal Periodontal Disease and Selected Adverse Pregnancy Outcomes

Pregnancy outcome	Findings from published research	Interpretation of evidence	Clinical consideration
Preterm birth	Several observational studies report increased risk, while others show no association	Inconsistent association; causation unproven	Assess obstetric risk independently of oral findings
Low birth weight	Observational research and reviews report associations in some populations	Potential association affected by study differences	Promote oral healthcare without promising neonatal benefit
Preeclampsia	Several studies report increased odds among women with periodontitis	Mixed evidence; possible shared inflammatory pathways	Continue established prenatal blood pressure monitoring
Fetal growth restriction	Some studies investigate associations with periodontal disease	Limited and inconsistent evidence	Use established obstetric growth surveillance
Small-for-gestational-age birth	Associations have been examined in prospective studies	Relationship remains uncertain	Evaluate maternal and placental risk factors
Pregnancy loss and stillbirth	Investigated as potential outcomes in some studies	Insufficient evidence for a direct causal	Follow established obstetric evaluation

		relationship	protocols
Periodontal inflammation	Treatment generally improves local periodontal indicators	Direct oral health benefit is established	Provide indicated dental treatment during pregnancy

Maternal Periodontal Disease, Preeclampsia, and Fetal Growth-Related Complications

Maternal periodontal disease has also been investigated in relation to preeclampsia, fetal growth restriction, and other pregnancy complications. These outcomes differ in their biological characteristics but may involve maternal inflammation, vascular abnormalities, metabolic disturbances, and altered placental function. Researchers have proposed that chronic periodontal inflammation could contribute to these processes through inflammatory mediators or microbial exposure. However, the available evidence remains heterogeneous, and direct causal relationships have not been established (Sanz & Kornman, 2013). Preeclampsia is a potentially serious hypertensive disorder of pregnancy that can affect maternal kidneys, liver, nervous system, and other organs. Abnormal placental development and systemic endothelial dysfunction play important roles in the condition. Because periodontal disease also involves chronic inflammatory activity, investigators have explored whether maternal periodontitis is associated with an increased risk of preeclampsia. A prospective study reported a relationship between periodontal disease and subsequent preeclampsia, contributing to the hypothesis that chronic maternal oral infection might be associated with obstetric vascular complications (Boggess et al., 2003). A case-control investigation conducted in Brazil further examined the association between maternal periodontitis and preeclampsia. The researchers identified a statistically significant relationship between periodontal disease and preeclampsia after considering selected maternal characteristics. These findings suggested that periodontal infection might contribute to the overall inflammatory burden experienced by pregnant women. However, the observational design could not establish whether periodontal disease preceded the pathological processes responsible for preeclampsia or whether both conditions reflected shared risk factors (Cota et al., 2006). Not all investigations have supported a consistent association. A multicenter prospective cohort study assessed periodontal disease in early pregnancy and examined preeclampsia, preterm birth, fetal growth restriction, and perinatal death. After adjustment for relevant confounding variables, periodontal disease was not associated with the combined adverse pregnancy outcome or with preeclampsia individually. Such findings emphasize that the relationship between oral health and hypertensive pregnancy disorders may vary across populations and may be influenced by differences in periodontal disease classification, maternal characteristics, and study methodology (Srinivas et al., 2009). Fetal growth restriction represents another important obstetric outcome that has been examined in relation to periodontal health. Inadequate fetal growth can result from maternal disease, placental dysfunction, nutritional factors, and other pregnancy-related complications. Researchers have hypothesized that chronic inflammation associated with periodontal infection could potentially influence placental vascular function or nutrient exchange. Nevertheless, these proposed mechanisms remain under investigation, and the available evidence does not establish periodontal disease as an independent cause of fetal growth restriction (Xiong et al., 2007). A prospective investigation examined maternal periodontal disease during early pregnancy and the risk of delivering a small-for-gestational-age infant. The findings suggested an association between periodontal disease and fetal growth-related outcomes in the studied population. However, small-for-gestational-age birth is influenced by multiple maternal and fetal characteristics, making careful adjustment for confounding essential. Consequently, the findings should be considered evidence of a possible epidemiological relationship rather than proof that periodontal infection directly restricts fetal growth (Boggess et al., 2006). Maternal metabolic conditions may further complicate the relationship between periodontal health and pregnancy outcomes. Diabetes and altered glucose regulation can influence inflammatory responses and increase susceptibility to periodontal disease. Gestational diabetes is also associated with adverse maternal and neonatal outcomes, creating potential overlap between oral and systemic health risks. Researchers have investigated whether periodontal disease is associated with gestational

diabetes, but the direction and independence of this relationship remain uncertain. Shared risk factors, including obesity, metabolic dysfunction, and socioeconomic circumstances, should be carefully considered (Xiong et al., 2007). Pregnancy loss and stillbirth have also been investigated as potential outcomes associated with maternal periodontal disease. These events may result from chromosomal abnormalities, maternal medical conditions, placental complications, infections, and other factors. Although some observational investigations have included pregnancy loss among their outcomes, available evidence is insufficient to conclude that periodontal infection independently causes miscarriage or stillbirth. Interpretation requires particular caution because these outcomes are relatively uncommon and may involve multiple overlapping clinical mechanisms (Xiong et al., 2006). A broader assessment of maternal oral health should therefore recognize both the biological plausibility of systemic inflammatory relationships and the limitations of available epidemiological evidence. Periodontal disease may occur alongside other conditions associated with complicated pregnancies, but it should not be treated as a substitute marker for established obstetric risk assessment. Pregnant women require appropriate prenatal monitoring regardless of their periodontal status, while dental disease should be evaluated and treated according to clinical need (Sanz & Kornman, 2013).

Preventive Oral Healthcare, Periodontal Treatment, and Maternal Health Outcomes

Preventive oral healthcare is an important component of maternal health because pregnancy-related changes may increase susceptibility to gingival inflammation and complicate the management of existing dental conditions. Appropriate prevention includes regular toothbrushing with fluoride toothpaste, interdental cleaning, dietary counseling, and professional dental assessment when indicated. These measures support periodontal health, reduce dental discomfort, and promote oral hygiene during pregnancy. However, the established benefits of oral healthcare should be distinguished from the less certain effects of dental treatment on obstetric outcomes (Silk et al., 2008).

Professional periodontal treatment generally involves examination, oral hygiene instruction, and removal of plaque and calculus through appropriate cleaning procedures. Patients with periodontitis may require scaling and root planing to reduce periodontal inflammation and improve clinical attachment-related indicators. Treatment plans should consider disease severity, symptoms, gestational stage, and relevant maternal medical conditions. The purpose of periodontal therapy is to control local infection and preserve oral health, regardless of whether it influences the risk of preterm birth or other pregnancy complications (Boggess & Edelstein, 2006). The potential effect of periodontal treatment on birth outcomes has been examined through randomized clinical trials. An early intervention study conducted in Chile investigated whether treatment of periodontal disease during pregnancy could reduce preterm low birth weight. The investigators reported findings suggesting a reduction in adverse birth outcomes following periodontal therapy. Although the results contributed to interest in prenatal periodontal intervention, subsequent trials produced less consistent findings. Differences in patient populations, disease definitions, treatment approaches, and baseline obstetric risks may explain some of the variation (López et al., 2002). A major randomized controlled trial evaluated nonsurgical periodontal treatment in pregnant women with periodontitis. The investigators assigned participants to receive periodontal treatment during pregnancy or after delivery. Periodontal therapy improved maternal periodontal health but did not significantly reduce preterm birth. Preterm delivery occurred in approximately 12.0% of treated participants and 12.8% of controls with available pregnancy outcome information. These findings demonstrate that treatment of periodontal disease can improve oral health without necessarily changing obstetric outcomes (Michalowicz et al., 2006). New England Journal of Medicine A separate randomized controlled trial in Australia investigated whether periodontal treatment during pregnancy could prevent premature delivery or other complications. The study similarly found that periodontal therapy improved oral health but did not significantly reduce preterm birth or other major adverse pregnancy outcomes. These findings provide important evidence against assuming that improvement in periodontal inflammation automatically produces measurable obstetric benefits. They also reinforce the need to distinguish clinical effectiveness for oral disease from effectiveness in preventing pregnancy complications (Newnham et al., 2009). Systematic reviews have synthesized the findings of

randomized trials evaluating periodontal treatment during pregnancy. A Cochrane review including 15 studies and 7,161 participants found no clear evidence that periodontal treatment reduced preterm birth before 37 weeks. The review identified low-certainty evidence suggesting that treatment might reduce low birth weight, but confidence in this estimate was limited. The authors concluded that further research was necessary to determine whether periodontal interventions could reliably prevent adverse pregnancy outcomes (Iheozor-Ejiofor et al., 2017). Cochrane A meta-analysis of randomized controlled trials also examined the relationship between periodontal therapy and birth outcomes. The investigators evaluated whether treating periodontal disease during pregnancy could influence the frequency of preterm birth and low birth weight. Although some individual trials reported favorable results, the combined evidence was insufficient to demonstrate consistent protection across all pregnancy outcomes. This variation highlights the importance of considering study quality, treatment timing, periodontal disease severity, and baseline maternal risk when interpreting intervention findings (George et al., 2011). Clinical recommendations should therefore avoid presenting periodontal treatment as a proven method for preventing premature delivery or preeclampsia. Nevertheless, dental assessment and necessary treatment remain appropriate during pregnancy because untreated oral disease can cause pain, infection, impaired chewing, and reduced quality of life. Healthcare professionals should explain that periodontal treatment improves oral health while the evidence concerning prevention of obstetric complications remains uncertain (Michalowicz et al., 2006). Prenatal oral health education is particularly valuable for women who have limited access to dental services or who are unfamiliar with pregnancy-related oral health changes. Education should address proper toothbrushing, interdental cleaning, prevention of dental caries, and the importance of seeking care for persistent bleeding, tooth pain, swelling, or signs of infection. Obstetric professionals can contribute by asking about dental symptoms and encouraging appropriate dental consultation. Clear communication can help reduce unnecessary postponement of treatment during pregnancy (Silk et al., 2008). Interdisciplinary care provides additional opportunities to strengthen maternal health services. Obstetricians, midwives, nurses, dentists, and dental hygienists can collaborate to identify oral health concerns and coordinate appropriate treatment. Such collaboration is particularly important for women with diabetes, significant periodontal disease, poor oral hygiene, or limited access to healthcare. Integrating oral health into prenatal assessment may improve service accessibility and reduce fragmentation between dental and maternal healthcare systems (Boggess & Edelstein, 2006).


Future Directions and Evidence-Based Strategies for Improving Maternal Oral Health During Pregnancy

Future research concerning maternal oral health and pregnancy outcomes should prioritize distinguishing epidemiological associations from causal relationships. Although numerous observational investigations have reported links between periodontal disease and complications such as preterm birth, low birth weight, and preeclampsia, findings remain inconsistent across different populations. Randomized trials have generally demonstrated improvements in periodontal health without consistently reducing adverse obstetric outcomes. Future investigations should therefore examine biological mechanisms, potential confounding factors, and clinically meaningful interventions using rigorous research designs (Sanz & Kornman, 2013). Standardization of periodontal disease assessment is an important research priority. Previous investigations have used different combinations of probing depth, clinical attachment loss, bleeding, and other periodontal indicators to classify participants. These differences may contribute to variation in reported associations and make comparisons between studies difficult. Future research should use clearly defined periodontal measurements, report disease severity, and distinguish gingivitis from destructive periodontitis. Standardized diagnostic criteria would strengthen the reliability of evidence synthesis and improve the interpretation of maternal oral health risks (Xiong et al., 2006). Longitudinal investigations beginning before conception may provide valuable information about the timing and development of periodontal inflammation. Many existing studies assess oral health after pregnancy has already begun, making it difficult to determine whether periodontal disease preceded early placental or inflammatory

changes. Preconception cohorts could examine how chronic oral conditions, maternal metabolic health, and pregnancy-related immune changes interact over time. Such research may help clarify whether periodontal disease contributes independently to pregnancy risk or primarily reflects shared maternal health factors (Boggess & Edelstein, 2006). Further investigation of biological markers may also improve understanding of possible oral-systemic relationships. Researchers could examine inflammatory mediators, periodontal microbial profiles, maternal immune responses, and placental indicators across different stages of pregnancy. Such investigations should consider whether measured biological differences are specific to periodontal disease or occur in association with obesity, smoking, diabetes, and other inflammatory conditions. Establishing biological plausibility is important, but a mechanistic finding alone would not demonstrate that periodontal disease causes adverse pregnancy outcomes (Sanz & Kornman, 2013).

Future intervention trials should evaluate whether particular groups of pregnant women benefit more from periodontal treatment. Women with severe periodontitis, high inflammatory burden, or specific obstetric risk profiles may differ from those with mild oral inflammation. However, subgroup effects should be investigated prospectively rather than inferred from small secondary analyses. Studies should also consider the timing of treatment, consistency of intervention delivery, and changes in periodontal health after treatment. Such approaches may help explain why earlier clinical trials produced different findings (Newnham et al., 2009). Research evaluating periodontal interventions should include both oral health and obstetric outcomes. Improvements in gingival bleeding, probing depth, and periodontal inflammation are clinically meaningful even when birth outcomes remain unchanged. At the same time, investigations designed to assess preterm birth or low birth weight should include adequate sample sizes and appropriate adjustment for established obstetric risk factors. Evidence syntheses have emphasized the need for consistent outcome reporting and sufficiently rigorous trials to determine whether treatment affects pregnancy complications (Iheozor-Ejiofor et al., 2017). Maternal oral health education should become a more consistent component of prenatal healthcare. Pregnancy provides an opportunity for healthcare professionals to discuss oral hygiene, diet, dental symptoms, and access to preventive services. Educational initiatives may be particularly valuable for first-time mothers and women who have not received routine dental care. However, education programs should be evaluated according to actual changes in oral hygiene practices, dental attendance, untreated disease, and patient knowledge rather than assuming that information provision alone improves health outcomes (Silk et al., 2008). Healthcare access should be addressed as an important determinant of maternal oral health. Financial limitations, geographical barriers, limited dental insurance, and misconceptions concerning treatment safety may prevent pregnant women from obtaining professional care. These barriers can contribute to untreated periodontal disease and dental infections regardless of whether oral conditions independently influence obstetric outcomes. Public health initiatives should therefore prioritize equitable access to preventive and therapeutic dental services as part of comprehensive maternal healthcare (Boggess & Edelstein, 2006). Interprofessional education may improve communication between dental and obstetric professionals. Obstetric clinicians should understand common oral health changes during pregnancy and recognize when dental assessment is indicated. Dental professionals should be familiar with relevant pregnancy-related precautions and communicate appropriately with maternal healthcare providers when complex medical conditions are present. Joint educational initiatives, standardized referral procedures, and coordinated clinical records may improve continuity of care and reduce unnecessary delays in treatment (Silk et al., 2008). Digital health technologies may offer opportunities to improve maternal oral health awareness and referral coordination. Electronic prenatal records could include basic oral health screening questions, documented dental referrals, and follow-up information. Mobile health applications may support oral hygiene reminders and provide access to appropriate educational resources. However, digital interventions should be designed to complement direct dental assessment rather than replace professional examination. Their effectiveness should be evaluated through measurable changes in oral health outcomes and service access (Boggess & Edelstein, 2006). International research collaboration may help explain why

associations between periodontal disease and pregnancy outcomes differ across populations. Differences in nutrition, socioeconomic circumstances, oral hygiene practices, dental healthcare availability, smoking exposure, and obstetric services may influence study findings. Comparative research should evaluate these factors systematically and determine whether apparent geographical differences persist after applying standardized periodontal and pregnancy outcome definitions. This approach could strengthen the external validity of future investigations (Xiong et al., 2007). Future evidence synthesis should place greater emphasis on the methodological quality of observational and intervention studies. Associations should be evaluated according to the likelihood of residual confounding, selection bias, measurement error, and inconsistent diagnostic criteria. Randomized trials should be assessed separately from observational studies because they answer different research questions. For example, observational studies investigate whether periodontal disease is associated with adverse pregnancy outcomes, whereas treatment trials evaluate whether a particular intervention changes those outcomes. These distinctions are essential for drawing clinically appropriate conclusions (Sanz & Kornman, 2013). The development of evidence-based maternal oral healthcare policies should prioritize established dental health needs while remaining responsive to emerging research. Pregnant women should be encouraged to maintain oral hygiene, seek professional assessment for symptoms, and receive appropriate treatment for existing dental conditions. Healthcare policies should not require proof of obstetric benefit before supporting access to necessary oral healthcare. Preventing dental pain, infection, and periodontal deterioration represents a sufficient clinical justification for appropriate dental services during pregnancy (Michalowicz et al., 2006). Ultimately, future progress will depend on integrating oral healthcare into broader maternal health strategies while maintaining scientific caution concerning proposed relationships with pregnancy complications. Continued collaboration between researchers, dentists, obstetricians, nurses, and public health professionals can strengthen understanding of maternal oral-systemic health. Evidence-based prevention, equitable healthcare access, and carefully designed research may improve maternal oral health and clarify whether specific interventions influence adverse pregnancy outcomes (George et al., 2011).



Summary:

Maternal oral health is an important component of prenatal healthcare because pregnancy-related physiological changes can influence gingival inflammation, periodontal conditions, and overall dental well-being. Periodontal disease has received particular attention because of its possible association

with adverse pregnancy outcomes, including preterm birth, low birth weight, preeclampsia, and fetal growth restriction. Observational studies have identified associations between maternal periodontal disease and certain adverse obstetric outcomes in some populations. However, findings are inconsistent, and the influence of shared maternal risk factors remains an important consideration. Proposed mechanisms involving systemic inflammation, microbial exposure, and placental dysfunction provide biological plausibility but do not establish causation. Randomized clinical trials demonstrate that periodontal treatment during pregnancy can improve maternal periodontal health. Nevertheless, treatment has not consistently reduced preterm birth or other major pregnancy complications. These findings emphasize the importance of distinguishing the established benefits of dental treatment from unproven obstetric effects. Comprehensive prenatal healthcare should include oral health education, appropriate dental assessment, timely treatment of existing oral disease, and effective communication between dental and obstetric professionals. Future investigations should use standardized periodontal definitions, rigorous study designs, and clearly defined pregnancy outcomes. Improving maternal oral healthcare remains clinically valuable because it can reduce untreated dental disease, improve comfort, and support general maternal well-being. A coordinated, evidence-based approach may strengthen the quality and accessibility of prenatal dental services while supporting continued research into the relationship between oral health and pregnancy outcomes.

References:

Boggess, K. A., Beck, J. D., Murtha, A. P., Moss, K., & Offenbacher, S. (2006). Maternal periodontal disease in early pregnancy and risk for a small-for-gestational-age infant. American Journal of Obstetrics and Gynecology, 194(5), 1316–1322. https://doi.org/10.1016/j.ajog.2005.11.059

Boggess, K. A., & Edelstein, B. L. (2006). Oral health in women during preconception and pregnancy: Implications for birth outcomes and infant oral health. Maternal and Child Health Journal, 10(5 Suppl.), S169–S174. https://doi.org/10.1007/s10995-006-0095-x

Cota, L. O. M., Guimarães, A. N., Costa, J. E., Lorentz, T. C. M., & Costa, F. O. (2006). Association between maternal periodontitis and an increased risk of preeclampsia. Journal of Periodontology, 77(12), 2063–2069. https://doi.org/10.1902/jop.2006.060061

George, A., Shamim, S., Johnson, M., Ajwani, S., Bhole, S., Blinkhorn, A., Ellis, S., & Andrews, K. (2011). Periodontal treatment during pregnancy and birth outcomes: A meta-analysis of randomised trials. International Journal of Evidence-Based Healthcare, 9(2), 122–147. https://doi.org/10.1111/j.1744-1609.2011.00210.x

Iheozor-Ejiofor, Z., Middleton, P., Esposito, M., & Glenny, A.-M. (2017). Treating periodontal disease for preventing adverse birth outcomes in pregnant women. Cochrane Database of Systematic Reviews, 2017(6), CD005297. https://doi.org/10.1002/14651858.CD005297.pub3

Jeffcoat, M. K., Geurs, N. C., Reddy, M. S., Cliver, S. P., Goldenberg, R. L., & Hauth, J. C. (2001). Periodontal infection and preterm birth: Results of a prospective study. The Journal of the American Dental Association, 132(7), 875–880. https://doi.org/10.14219/jada.archive.2001.0299

López, N. J., Smith, P. C., & Gutierrez, J. (2002). Periodontal therapy may reduce the risk of preterm low birth weight in women with periodontal disease: A randomized controlled trial. Journal of Periodontology, 73(8), 911–924. https://pubmed.ncbi.nlm.nih.gov/12211502/

Michalowicz, B. S., Hodges, J. S., DiAngelis, A. J., Lupo, V. R., Novak, M. J., Ferguson, J. E., Buchanan, W., Bofill, J., Papapanou, P. N., Mitchell, D. A., Matseoane, S., & Tschida, P. A. (2006). Treatment of periodontal disease and the risk of preterm birth. The New England Journal of Medicine, 355(18), 1885–1894. https://doi.org/10.1056/NEJMoa062249

Newnham, J. P., Newnham, I. A., Ball, C. M., Wright, M., Pennell, C. E., Swain, J., & Doherty, D. A. (2009). Treatment of periodontal disease during pregnancy: A randomized controlled trial. Obstetrics & Gynecology, 114(6), 1239–1248. https://doi.org/10.1097/AOG.0b013e3181c15b40

Offenbacher, S., Katz, V., Fertik, G., Collins, J., Boyd, D., Maynor, G., McKaig, R., & Beck, J. (1996). Periodontal infection as a possible risk factor for preterm low birth weight. Journal of Periodontology, 67(10 Suppl.), 1103–1113. https://doi.org/10.1902/jop.1996.67.10s.1103

Sanz, M., & Kornman, K. (2013). Periodontitis and adverse pregnancy outcomes: Consensus report of the Joint EFP/AAP Workshop on Periodontitis and Systemic Diseases. Journal of Clinical Periodontology, 40(Suppl. 14), S164–S169. https://doi.org/10.1111/jcpe.12083

Silk, H., Douglass, A. B., Douglass, J. M., & Silk, L. (2008). Oral health during pregnancy. American Family Physician, 77(8), 1139–1144. https://www.aafp.org/afp/2008/0415/p1139

Srinivas, S. K., Sammel, M. D., Stamilio, D. M., Clothier, B., Jeffcoat, M. K., Parry, S., Macones, G. A., Elovitz, M. A., & Metlay, J. (2009). Periodontal disease and adverse pregnancy outcomes: Is there an association? American Journal of Obstetrics and Gynecology, 200(5), 497.e1–497.e8. https://doi.org/10.1016/j.ajog.2009.03.003

Xiong, X., Buekens, P., Fraser, W. D., Beck, J., & Offenbacher, S. (2006). Periodontal disease and adverse pregnancy outcomes: A systematic review. BJOG: An International Journal of Obstetrics & Gynaecology, 113(2), 135–143. https://doi.org/10.1111/j.1471-0528.2005.00827.x

Xiong, X., Buekens, P., Vastardis, S., & Yu, S. M. (2007). Periodontal disease and pregnancy outcomes: State-of-the-science. Obstetrical & Gynecological Survey, 62(9), 605–615. https://doi.org/10.1097/01.ogx.0000279292.63435.40

Hartnett, E., Haber, J., Krainovich-Miller, B., Bella, A., Vasilyeva, A., & Kessler, J. L. (2016). Oral health in pregnancy. Journal of Obstetric, Gynecologic & Neonatal Nursing, 45(4), 565–573. https://doi.org/10.1016/j.jogn.2016.04.005

Figuero, E., Carrillo-de-Albornoz, A., Herrera, D., & Bascones-Martínez, A. (2010). Gingival changes during pregnancy: I. Influence of hormonal variations on clinical and immunological parameters. Journal of Clinical Periodontology, 37(3), 220–229. https://doi.org/10.1111/j.1600-051X.2009.01516.x

Chen, P., Hong, F., & Yu, X. (2022). Prevalence of periodontal disease in pregnancy: A systematic review and meta-analysis. Journal of Dentistry, 125, 104253. https://doi.org/10.1016/j.jdent.2022.104253

Bett, J. V. S., Batistella, E. Â., Melo, G., Munhoz, E. de A., Silva, C. A. B., Guerra, E. N. da S., Porporatti, A. L., & Canto, G. De L. (2019). Prevalence of oral mucosal disorders during pregnancy: A systematic review and meta-analysis. Journal of Oral Pathology & Medicine, 48(4), 270–277. https://doi.org/10.1111/jop.12831

Offenbacher, S., Katz, V., Fertik, G., Collins, J., Boyd, D., Maynor, G., McKaig, R., & Beck, J. (1996). Periodontal infection as a possible risk factor for preterm low birth weight. Journal of Periodontology, 67(10 Suppl.), 1103–1113. https://doi.org/10.1902/jop.1996.67.10s.1103

Han, Y. W., Fardini, Y., Chen, C., Iacampo, K. G., Peraino, V. A., Shamonki, J. M., & Redline, R. W. (2010). Term stillbirth caused by oral Fusobacterium nucleatum. Obstetrics & Gynecology, 115(2 Pt 2), 442–445. https://doi.org/10.1097/AOG.0b013e3181cb9955

Bright, R., Macowan, M. G., Tian, K., Fitzsimmons, T., Wilson, R. L., Roberts, C. T., Christophersen, C. T., Bartold, P. M., Kidd, S. P., & Zilm, P. S. (2026). Periodontitis during pregnancy: The effect on the gut microbiome and intestinal inflammation. Journal of Periodontology, 97(7), 1587–1603. https://doi.org/10.1002/jper.70132

Valentine, G. C., Bobetsis, Y. A., Shayo, B. C., Milgrom, P., & Madianos, P. N. (2026). Periodontitis and adverse pregnancy outcomes: Mechanistic evidence. Journal of Periodontal Research. Advance online publication. https://doi.org/10.1111/jre.70134

Manrique-Corredor, E. J., Orozco-Beltran, D., Lopez-Pineda, A., Quesada, J. A., Gil-Guillen, V. F., & Carratala-Munuera, C. (2019). Maternal periodontitis and preterm birth: Systematic review and meta-analysis. Community Dentistry and Oral Epidemiology, 47(3), 243–251. https://doi.org/10.1111/cdoe.12450

Khader, Y. S., & Ta'ani, Q. (2005). Periodontal diseases and the risk of preterm birth and low birth weight: A meta-analysis. Journal of Periodontology, 76(2), 161–165. https://doi.org/10.1902/jop.2005.76.2.161

Corbella, S., Taschieri, S., Del Fabbro, M., Francetti, L., Weinstein, R., & Ferrazzi, E. (2016). Adverse pregnancy outcomes and periodontitis: A systematic review and meta-analysis exploring

potential association. Quintessence International, 47(3), 193–204. https://doi.org/10.3290/j.qi.a34980

Montoya-Carralero, J.-M., Ávila-Villasmil, R., Sánchez-Pérez, A., Jornet-García, A., Terrer-Alonso, E., & Moya-Villaescusa, M.-J. (2024). Relationship between periodontal disease and preterm birth. A systematic review and meta-analysis. Medicina Oral, Patología Oral y Cirugía Bucal, 29(6), e857–e865. https://doi.org/10.4317/medoral.26830

Le, Q.-A., Akhter, R., Coulton, K. M., Vo, N. T. N., Duong, L. T. Y., Nong, H. V., Yaacoub, A., Condous, G., Eberhard, J., & Nanan, R. (2022). Periodontitis and preeclampsia in pregnancy: A systematic review and meta-analysis. Maternal and Child Health Journal, 26(12), 2419–2443. https://doi.org/10.1007/s10995-022-03556-6

Huang, X., Wang, J., Liu, J., Hua, L., Zhang, D., Hu, T., & Ge, Z.-L. (2014). Maternal periodontal disease and risk of preeclampsia: A meta-analysis. Journal of Huazhong University of Science and Technology [Medical Sciences], 34(5), 729–735. https://doi.org/10.1007/s11596-014-1343-8

Boggess, K. A., Beck, J. D., Murtha, A. P., Moss, K., & Offenbacher, S. (2006). Maternal periodontal disease in early pregnancy and risk for a small-for-gestational-age infant. American Journal of Obstetrics and Gynecology, 194(5), 1316–1322. https://doi.org/10.1016/j.ajog.2005.11.059

Gare, J., Kanoute, A., Meda, N., Viennot, S., Bourgeois, D., & Carrouel, F. (2021). Periodontal conditions and pathogens associated with pre-eclampsia: A scoping review. International Journal of Environmental Research and Public Health, 18(13), 7194. https://doi.org/10.3390/ijerph18137194

Michalowicz, B. S., Hodges, J. S., DiAngelis, A. J., Lupo, V. R., Novak, M. J., Ferguson, J. E., Buchanan, W., Bofill, J., Papapanou, P. N., Mitchell, D. A., Matseoane, S., & Tschida, P. A. (2006). Treatment of periodontal disease and the risk of preterm birth. The New England Journal of Medicine, 355(18), 1885–1894. https://doi.org/10.1056/NEJMoa062249

Newnham, J. P., Newnham, I. A., Ball, C. M., Wright, M., Pennell, C. E., Swain, J., & Doherty, D. A. (2009). Treatment of periodontal disease during pregnancy: A randomized controlled trial. Obstetrics & Gynecology, 114(6), 1239–1248. https://doi.org/10.1097/AOG.0b013e3181c15b40

Offenbacher, S., Beck, J. D., Jared, H. L., Mauriello, S. M., Mendoza, L. C., Couper, D. J., Stewart, D. D., Murtha, A. P., Cochran, D. L., Dudley, D. J., Reddy, M. S., Geurs, N. C., & Hauth, J. C. (2009). Effects of periodontal therapy on rate of preterm delivery: A randomized controlled trial. Obstetrics & Gynecology, 114(3), 551–559. https://doi.org/10.1097/AOG.0b013e3181b1341f

Bao, J., Huang, X., Wang, L., He, Y., Rasubala, L., & Ren, Y.-F. (2022). Clinical practice guidelines for oral health care during pregnancy: A systematic evaluation and summary recommendations for general dental practitioners. Quintessence International, 53(4), 362–373. https://doi.org/10.3290/j.qi.b2644863

Sanz, M., & Kornman, K. (2013). Periodontitis and adverse pregnancy outcomes: Consensus report of the Joint EFP/AAP Workshop on Periodontitis and Systemic Diseases. Journal of Clinical Periodontology, 40(Suppl. 14), S164–S169. https://doi.org/10.1111/jcpe.12083

Xu, H., Cai, M., Xu, H., Shen, X.-J., & Liu, J. (2025). Role of periodontal treatment in pregnancy gingivitis and adverse outcomes: A systematic review and meta-analysis. The Journal of Maternal-Fetal & Neonatal Medicine, 38(1), 2416595. https://doi.org/10.1080/14767058.2024.2416595

Da Rosa, M. I., Pires, P. D. S., Medeiros, L. R., Edelweiss, M. I., & Martínez-Mesa, J. (2012). Periodontal disease treatment and risk of preterm birth: A systematic review and meta-analysis. Cadernos de Saúde Pública, 28(10), 1823–1833. https://doi.org/10.1590/S0102-311X2012001000002

Quemeneur, A., & Boutigny-Vella, H. (2026). Periodontal diseases and pregnancy. European Journal of Obstetrics & Gynecology and Reproductive Biology, 327, 115425. https://doi.org/10.1016/j.ejogrb.2026.115425