Birat Journal Of Health Sciences
ISSN (Online): 2542-2804 • Quarterly • Vol. 8, No. 2 (2022)

Evaluation of Pediatric Emergency Admission Patterns and Seasonal Variations in Disease Presentation

Reema Harrison[1]
[1] School of Population Health, Faculty of Medicine and Health, UNSW Sydney, Australia
Email: reema.harrison@unsw.edu.au

ARTICLE HISTORY

RECEIVED	REVISED	ACCEPTED	PUBLISHED ONLINE
[15 May 2022]	*[18 July 2022]*	*[23 August 2022]*	*[31 August 2022]*

Abstract

Pediatric emergency departments play an essential role in providing immediate medical assessment and treatment for children presenting with acute illnesses, injuries, and potentially life-threatening conditions. Understanding pediatric emergency admission patterns and seasonal variations in disease presentation is important for improving healthcare resource allocation, clinical preparedness, and patient outcomes. This narrative review evaluates the demographic, clinical, environmental, and organizational factors influencing pediatric emergency utilization, with particular emphasis on seasonal variations in respiratory infections, gastrointestinal disorders, asthma exacerbations, traumatic injuries, and other common childhood conditions. Evidence from observational studies and epidemiological investigations indicates that pediatric emergency visits frequently demonstrate predictable temporal patterns influenced by age, infectious disease circulation, environmental conditions, school calendars, and healthcare accessibility. Respiratory illnesses commonly contribute to increased emergency visits during colder months in temperate regions, while gastrointestinal disorders, allergic conditions, and injuries may demonstrate different seasonal distributions. Nevertheless, seasonal patterns vary considerably across geographical regions, climatic conditions, and patient populations. Variations in emergency attendance, hospital admission, and disease severity require separate evaluation because increased patient volume does not necessarily indicate a corresponding rise in serious illness. The review highlights the importance of systematic disease surveillance, age-specific clinical assessment, and forecasting models for improving pediatric emergency service planning. Understanding these patterns can support targeted prevention, effective staffing, and timely emergency care for children throughout the year.

Keywords: Pediatric Emergency Department, Seasonal Variation, Emergency Admissions, Childhood Diseases, Respiratory Infections, Gastrointestinal Disorders, Pediatric Asthma, Disease Epidemiology, Emergency Healthcare Utilization, Healthcare Resource Planning

COPYRIGHT & LICENSE

Copyright © 2026 The Author(s). All rights reserved by the author(s), who retain copyright ownership of their original work. The author(s) grant the **Birat Journal of Health Sciences (BJHS)** a non-exclusive right of first publication.

Open Access License: This article is published under the [Creative Commons Attribution 4.0 International (CC BY 4.0) License](https://creativecommons.org/licenses/by/4.0/), permitting unrestricted use, distribution, reproduction, and adaptation in any medium, including commercial use, provided that appropriate credit is given to the original author(s), the source and the license, and any modifications are clearly indicated.

1. Introduction

Pediatric emergency departments provide essential healthcare services for infants, children, and adolescents experiencing acute illnesses, traumatic injuries, and medical conditions requiring urgent evaluation. These services frequently manage a broad spectrum of clinical presentations, ranging from

uncomplicated respiratory infections and minor injuries to severe dehydration, respiratory distress, neurological emergencies, and serious systemic infections. Effective pediatric emergency care requires sufficient clinical resources, timely triage, accurate diagnosis, and appropriate treatment according to the severity of each patient's condition. The number and characteristics of children presenting to emergency departments are not uniform throughout the year. Emergency utilization may change according to seasonal infectious disease activity, environmental conditions, school attendance patterns, public holidays, and population movement. Such variations can influence the demand for medical personnel, inpatient beds, diagnostic testing, and emergency treatment facilities. Seasonal changes in respiratory infections, asthma exacerbations, gastrointestinal illnesses, and certain environmental conditions may contribute to predictable fluctuations in pediatric healthcare utilization. However, the timing and magnitude of these fluctuations differ across geographical regions. Climatic conditions, socioeconomic circumstances, vaccination coverage, population density, and access to primary healthcare services may influence local patterns of emergency presentation. An important distinction exists between emergency department attendance and hospital admission. Many children receive assessment and treatment before being discharged home, while others require admission to pediatric wards or intensive care units. Evaluating total emergency visits without considering illness severity, triage category, and admission outcomes may therefore provide an incomplete understanding of pediatric emergency healthcare requirements. This article examines pediatric emergency admission patterns and seasonal variations in disease presentation through six interconnected areas: demographic and temporal characteristics, seasonal infectious diseases, respiratory and allergic conditions, gastrointestinal and injury-related presentations, healthcare service utilization, and future approaches to prediction and resource planning.

Demographic Characteristics and Temporal Patterns of Pediatric Emergency Admissions

Pediatric emergency utilization is influenced by demographic characteristics, disease severity, healthcare accessibility, and the timing of medical presentations. Children of different ages demonstrate distinct clinical needs because physiological development, immune function, mobility, and exposure to environmental hazards change throughout childhood. Infants and younger children frequently present with infectious illnesses, fever, and feeding-related concerns, while older children and adolescents may experience a broader distribution of traumatic injuries, behavioral conditions, and other acute medical problems. Understanding these differences is essential for evaluating emergency admission patterns and developing age-appropriate healthcare services (Gowa et al., 2019). A retrospective investigation conducted at the National Institute of Child Health in Karachi, Pakistan, examined the frequency and spectrum of illnesses presenting to a tertiary pediatric emergency department. Respiratory and gastrointestinal conditions represented major categories of clinical presentation, while patient attendance varied according to the time of day and day of the week. The study identified increased attendance during evening shifts and weekends, demonstrating that emergency department demand is affected by temporal patterns in addition to disease prevalence. These findings are relevant for healthcare institutions seeking to align staff availability with anticipated patient volume (Gowa et al., 2019). Age-related differences may also influence the proportion of children requiring inpatient treatment. Infants may need closer observation because of immature physiological regulation and the possibility of rapid deterioration during acute illness. Older children may present with conditions that can be managed through outpatient treatment, although the severity of individual presentations remains the primary determinant of admission. A Canadian study examining 46,942 pediatric emergency visits found that nighttime arrivals represented a smaller proportion of overall attendance but included a relatively greater proportion of high-acuity cases. This demonstrates that emergency volume and clinical severity should be considered separately when evaluating service requirements (Almubarak et al., 2019). Daily and weekly variations in attendance are particularly important for emergency department operations. Families may seek medical attention during evenings because of parental employment schedules, school attendance, or limited access to primary care outside regular working hours. Weekend visits may also increase when routine outpatient services are unavailable. In an investigation involving 223,677 pediatric emergency visits,

researchers observed higher daily attendance on Mondays and lower volumes during certain public holidays. The findings suggest that calendar-related patterns can provide useful information for predicting staffing needs and organizing pediatric emergency services (Krinsky-Diener et al., 2017). Population characteristics and geographical location can further affect emergency attendance. Hospitals located in tourist destinations or major referral centers may receive different patient groups during particular periods of the year. A study conducted in Charleston, South Carolina, found that local pediatric patients demonstrated increased emergency utilization from October through April, whereas out-of-state patients were more likely to attend during summer vacations and major holidays. These differences were associated with distinct clinical presentations and healthcare needs, indicating that population mobility can influence emergency demand independently of local disease seasonality (Rose et al., 2022). Seasonal variation should therefore be evaluated alongside age, sex, residence, socioeconomic circumstances, and healthcare access. Differences in emergency attendance may reflect changes in disease incidence, healthcare-seeking behavior, or the availability of alternative medical services. Hospital administrators should examine these factors when interpreting admission statistics and determining whether increased patient volume represents a genuine rise in severe illness or a change in healthcare utilization patterns (Solmaz & Orhan, 2026).

Seasonal Distribution of Infectious Diseases Among Pediatric Emergency Patients

Infectious diseases represent a substantial proportion of pediatric emergency presentations and contribute significantly to seasonal fluctuations in healthcare demand. Children frequently experience respiratory infections, gastrointestinal illnesses, ear infections, urinary tract infections, and other communicable conditions. The frequency of these illnesses may vary according to pathogen circulation, environmental temperature, humidity, school attendance, and patterns of social contact. Understanding these relationships is important for predicting emergency department workloads and preparing appropriate diagnostic and treatment resources (Lipsett et al., 2021). Respiratory infections are commonly associated with increased emergency attendance during colder months in temperate climates. Viral pathogens may spread more efficiently under certain environmental and behavioral conditions, while indoor crowding can increase opportunities for transmission. A national investigation of infectious disease-related pediatric emergency visits in the United States identified a pronounced winter increase across childhood age groups. The estimated number of infection-related visits was approximately 431,400 in July compared with 875,533 in February, demonstrating the considerable effect of seasonal infectious disease activity on emergency healthcare utilization (Hasegawa et al., 2015). The seasonal distribution of infectious illnesses differs according to the specific disease. A study examining common pediatric infectious conditions using a national emergency department database found seasonal patterns involving upper respiratory infections, pneumonia, otitis media, gastroenteritis, urinary tract infections, and skin infections. However, the strength and timing of seasonal variation differed between diagnostic categories. Pneumonia demonstrated a pronounced seasonal pattern, while other conditions exhibited more moderate fluctuations. These findings emphasize the importance of evaluating individual infectious diseases rather than assuming that all pediatric infections follow the same seasonal distribution (Lipsett et al., 2021). Infants represent a particularly important population because fever may indicate either a self-limited viral illness or a serious bacterial infection requiring urgent investigation. The seasonal circulation of viruses can influence the number of febrile infants presenting to emergency departments, but seasonal patterns should not replace clinical risk assessment. A retrospective study involving 1,529 febrile infants aged 90 days or younger found that viral infections demonstrated clear seasonal variation, including winter respiratory peaks and summer circulation of certain enteroviruses. In contrast, serious bacterial infections did not show the same seasonal pattern. These findings highlight the importance of maintaining appropriate bacterial infection assessment regardless of the time of year (Bellini et al., 2026). School calendars may also influence infectious disease transmission. Children attending schools and childcare facilities experience regular contact with peers, increasing opportunities for respiratory and gastrointestinal pathogen exposure. Periods following school reopening may therefore be associated with changes in healthcare demand. An eight-year study

involving 670,379 pediatric emergency visits in Lisbon identified recurring annual and shorter temporal cycles associated with respiratory infections, school calendars, and weather-related variables. The findings demonstrate that infectious disease patterns interact with broader social and environmental influences (Almeida et al., 2022). Changes in infectious disease presentation can also influence emergency department organization. During periods of high respiratory infection activity, hospitals may experience increased demand for clinical assessment, oxygen therapy, respiratory support, diagnostic testing, and infection prevention measures. However, resource requirements depend on the severity and age distribution of patients rather than the total number of visits alone. Seasonal surveillance should therefore include indicators of clinical acuity, hospitalization, and intensive care utilization in addition to the number of infectious disease presentations (Hasegawa et al., 2015). The evaluation of pediatric infectious disease seasonality should recognize geographical differences. In tropical and subtropical regions, respiratory and gastrointestinal diseases may follow patterns associated with rainfall, humidity, and local pathogen circulation rather than the four-season distribution observed in temperate climates. Healthcare organizations should therefore rely on local surveillance data and established regional epidemiological patterns when predicting disease-related emergency demand (Lipsett et al., 2021).

Respiratory Disorders, Asthma Exacerbations, and Seasonal Clinical Variations

Respiratory disorders are among the most important causes of pediatric emergency attendance because children may develop acute breathing difficulties associated with viral infections, asthma, bronchiolitis, pneumonia, and other respiratory conditions. Seasonal changes in respiratory disease presentation are influenced by infectious pathogens, environmental allergens, weather conditions, and age-related differences in airway physiology. Evaluating these factors can help emergency departments anticipate periods of increased respiratory care demand and prepare appropriate clinical resources (Dondi et al., 2017). Asthma is a major chronic respiratory condition associated with recurrent emergency visits among children. Acute exacerbations may occur following respiratory infections, allergen exposure, environmental pollution, and other triggers. However, the seasonal distribution of exacerbations varies according to age and geographical conditions. A retrospective study conducted in Italy found that respiratory infections were the principal triggers of acute asthma exacerbations among children, while allergic triggers were more prominent in school-aged patients. Preschool children demonstrated peaks during autumn and winter, whereas older children experienced different seasonal distributions (Dondi et al., 2017). Evidence from South Korea provides further insight into pediatric asthma seasonality. A nationwide investigation involving 41,128 children and adolescents presenting to emergency departments for asthma identified substantial seasonal differences. Asthma-related emergency visits generally increased during autumn and decreased during summer. The study also found variation across age categories and reported that a considerable proportion of asthma-related visits resulted in hospital admission. These findings demonstrate that environmental and temporal factors can affect both the frequency of presentations and the burden placed on pediatric inpatient services (Won et al., 2016). A six-year study conducted in Istanbul examined emergency visits among children with asthma between 2017 and 2022. The investigators found that asthma-related visits were concentrated more heavily during certain colder months, including October through January, with reduced attendance during summer. Younger children accounted for a substantial proportion of asthma presentations. These findings support the importance of regional surveillance and age-specific analysis when assessing seasonal respiratory disease patterns (Altaş et al., 2023). Respiratory disease patterns may also differ according to underlying diagnoses and the timing of infectious outbreaks. Viral respiratory infections can contribute to increased wheezing and respiratory distress among infants and preschool children, while older children may experience exacerbations related to seasonal allergens or environmental exposure. The use of broad diagnostic categories such as respiratory illness may therefore conceal clinically meaningful differences between asthma, bronchiolitis, pneumonia, and other respiratory conditions. More detailed diagnostic classification can support better prediction of equipment and treatment requirements (Lipsett et al., 2021). Recent research examining condition-specific seasonal

rhythms in pediatric emergencies identified widespread seasonal variation among different presenting complaints. Analysis of approximately 1.5 million pediatric emergency and urgent care encounters demonstrated that many conditions showed recognizable seasonal patterns, although peak timing differed considerably across diagnoses. Such findings suggest that healthcare institutions should move beyond forecasting total emergency attendance and examine the timing of individual disease categories. This approach may improve planning for respiratory therapy, diagnostic services, and specialized pediatric care (Tachinardi et al., 2026). Seasonal respiratory patterns have implications for preventive healthcare. Appropriate vaccination, asthma management education, medication adherence, and access to primary care may reduce certain preventable emergency presentations. Healthcare professionals should provide families with clear instructions concerning the recognition of respiratory distress and the circumstances requiring urgent evaluation. Preventive planning should also incorporate local infectious disease surveillance and environmental risk information rather than relying exclusively on historical monthly attendance patterns (Dondi et al., 2017). The clinical significance of respiratory seasonality extends beyond patient volume. Hospitals may experience increased demand for oxygen administration, nebulized treatment, respiratory monitoring, diagnostic imaging, and pediatric admission beds during periods of high disease activity. Appropriate staffing and equipment planning should therefore consider both the expected number of respiratory presentations and the likelihood of clinically significant illness. Integrating these indicators can support more responsive pediatric emergency care (Won et al., 2016).

Seasonal Patterns of Common Pediatric Emergency Presentations and Healthcare Planning Implications

Disease or presentation category	Common seasonal observation	Important contributing factors	Healthcare planning implications
Viral respiratory infections	Frequently increase during colder months in temperate regions	Viral circulation, indoor contact, school attendance	Additional respiratory assessment capacity
Bronchiolitis and infant respiratory illness	Seasonal peaks associated with local respiratory virus activity	Age, viral exposure, regional epidemiology	Respiratory monitoring and pediatric inpatient preparedness
Pediatric asthma exacerbations	Often increase during autumn or winter, depending on region	Respiratory infections, allergens, environmental exposures	Asthma treatment supplies and staff preparation
Pneumonia	Often demonstrates pronounced seasonal variation	Respiratory pathogen activity and patient susceptibility	Diagnostic testing and inpatient capacity planning
Acute gastroenteritis	Varies according to pathogen, climate, and location	Viral circulation, food and water exposure	Hydration assessment and infection prevention
Traumatic injuries	May increase during periods of outdoor activity or tourism	Recreation, school holidays, travel	Emergency procedures, imaging, and trauma resources
Febrile illness in young infants	Viral causes vary seasonally; bacterial risk persists throughout the year	Age, pathogen exposure, underlying clinical risk	Age-appropriate fever assessment protocols
Mental health emergencies	Some settings show spring and autumn peaks	School calendars and multiple social factors	Mental health assessment and referral capacity

Gastrointestinal Illnesses, Traumatic Injuries, and Other Seasonal Emergency Presentations

Gastrointestinal illnesses represent an important category of pediatric emergency presentations because vomiting, diarrhea, abdominal pain, and dehydration can require urgent medical evaluation. Acute gastroenteritis is particularly common in younger children, although its severity varies according to the causative organism, age, hydration status, and coexisting medical conditions. Seasonal patterns are influenced by pathogen circulation, food and water safety, environmental conditions, and community exposure. Consequently, gastrointestinal emergency attendance may increase at different times of the year depending on the geographical region and predominant infectious agents (Lipsett et al., 2021). The relationship between gastrointestinal infections and seasonal emergency demand has been demonstrated in population-based investigations. A United States study of common pediatric infectious diseases identified a significant seasonal pattern for gastroenteritis, although the magnitude of variation differed from respiratory conditions. These findings indicate that gastrointestinal disease contributes to predictable fluctuations in emergency utilization but should not be assumed to follow the same seasonal cycle as pneumonia or upper respiratory infections. Hospital surveillance systems should therefore analyze gastrointestinal presentations separately to improve forecasting and clinical preparedness (Lipsett et al., 2021). Regional studies provide additional evidence regarding the importance of gastrointestinal conditions. Research conducted in Karachi identified gastrointestinal emergencies alongside respiratory illnesses as major contributors to pediatric emergency workload. Such findings demonstrate that hospitals serving densely populated communities may encounter substantial demand for the management of vomiting, diarrhea, abdominal pain, and dehydration. However, individual hospital attendance patterns are influenced by local disease prevalence, healthcare accessibility, and referral arrangements. These factors should be considered before generalizing findings across different healthcare institutions (Gowa et al., 2019). The severity of gastrointestinal illness has important implications for admission decisions. Children with mild dehydration may respond to oral rehydration and observation, whereas those with severe dehydration, persistent vomiting, altered consciousness, or significant electrolyte abnormalities may require intravenous treatment or inpatient care. The number of gastrointestinal emergency visits does not therefore provide a complete measure of the demand for inpatient services. Evaluations should include clinical severity, treatment requirements, and admission outcomes in addition to diagnostic frequency (Solmaz & Orhan, 2026). Traumatic injuries represent another major category of pediatric emergency presentations. Falls, road traffic incidents, sports-related injuries, and recreational accidents may contribute to emergency attendance throughout childhood. The seasonal distribution of injuries can be influenced by outdoor activities, school holidays, tourism, weather conditions, and differences in supervision. In an investigation conducted in a destination city, out-of-state pediatric patients presented more frequently during summer and holiday periods and were more likely than local children to receive injury-related diagnoses. These findings illustrate how population movement and recreational activities can shape emergency healthcare utilization (Rose et al., 2022). Patterns of trauma may also vary according to time of day and school attendance. Children participating in organized recreation or outdoor activities may experience injuries during periods when such activities are common. A Canadian investigation found that minor trauma represented a leading category of daytime and evening emergency visits, whereas infectious conditions were relatively more prominent during nighttime attendance. This demonstrates that temporal variation within individual days can be as important as seasonal changes when planning emergency treatment services (Almubarak et al., 2019). Allergic conditions and skin-related presentations may demonstrate additional seasonal variation. Exposure to pollen, environmental allergens, insects, and other external factors can change according to local weather conditions and geographical characteristics. However, the relationship between these exposures and emergency attendance is complex because allergic symptoms may overlap with respiratory infections or other clinical conditions. In an eight-year analysis from Lisbon, peaks occurring outside the main winter respiratory season were associated with increased presentations among older children involving trauma, gastrointestinal illness, and atopic symptoms.

These findings support the value of disease-specific surveillance across multiple seasonal periods (Almeida et al., 2022). Mental health and behavioral emergencies also contribute to pediatric emergency utilization, particularly among older children and adolescents. Their temporal distribution may be influenced by school schedules, family circumstances, psychosocial stressors, and access to community mental health services. A recent analysis of approximately 1.5 million pediatric emergency and urgent care encounters identified substantial seasonal variation in psychiatric presentations, with peaks during spring and autumn. However, these patterns varied across geographical settings, and the study's observational design does not establish that seasonal factors directly caused the changes in attendance (Tachinardi et al., 2026). Comprehensive evaluation of nonrespiratory pediatric emergencies should therefore include gastrointestinal disorders, injuries, allergic conditions, neurological symptoms, and mental health presentations. Each category may follow distinct temporal patterns and require different clinical resources. Hospitals that examine only total patient attendance may fail to recognize changes in the specific services required during different seasons. Disease-specific analysis can support more effective planning without relying on overly simplified assumptions about seasonal illness (Guo et al., 2023).

Healthcare Service Utilization, Hospital Admission Rates, and Operational Challenges

Seasonal changes in pediatric emergency attendance can influence hospital operations by altering the demand for clinical assessments, diagnostic testing, inpatient beds, and specialist consultations. High-volume periods may create challenges involving waiting times, staff workload, treatment capacity, and patient flow. However, emergency department attendance and hospital admission represent distinct outcomes. A substantial increase in visits may be associated with common low-acuity illnesses, while a smaller number of severe presentations may require considerable inpatient resources. Healthcare organizations should therefore evaluate attendance patterns alongside triage severity and admission outcomes (Almubarak et al., 2019). Evidence from southeastern Türkiye illustrates the importance of distinguishing emergency visits from inpatient admission. A retrospective investigation involving 192,661 pediatric emergency visits during 2023 found that winter accounted for the largest proportion of attendance, followed by spring, autumn, and summer. Upper respiratory tract infections and acute gastroenteritis were the most frequently recorded diagnoses. Nevertheless, only approximately 0.94% of visits resulted in hospital admission, while 0.11% resulted in intensive care admission. These findings demonstrate that high patient volume may coexist with relatively low hospitalization rates, depending on the hospital's patient population and referral practices (Solmaz & Orhan, 2026). Seasonal variation in emergency utilization may also influence waiting times. When patient arrival rates increase beyond available clinical capacity, children and families may experience delays in registration, triage, medical evaluation, and discharge. A retrospective study using electronic medical record data from a pediatric emergency department in China found that patient attendance varied with school terms, holidays, and seasonal periods. Higher attendance was associated with longer waiting times, suggesting that predictable fluctuations in demand may provide opportunities for improved resource allocation (Guo et al., 2023). Temporal patterns can be examined through forecasting models that incorporate historical attendance data. An eight-year investigation of pediatric emergency visits in Lisbon identified annual, semiannual, weekly, and other recurring patterns. The researchers developed a time-series model incorporating school calendars and meteorological information, achieving a mean absolute percentage error of approximately 10.7% during cross-validation. Although prediction performance may differ in other settings, these findings suggest that emergency attendance can be forecast using a limited set of relevant temporal and environmental variables (Almeida et al., 2022). Long-term forecasting has also been evaluated in Australian pediatric emergency services. A study conducted at the Royal Children's Hospital in Melbourne used several years of attendance data to model seasonal variation in emergency presentations. The investigators identified recurring annual patterns characterized by lower attendance during summer and higher attendance during winter. The model explained a substantial proportion of historical variation, demonstrating the potential usefulness of quantitative methods for anticipating service demand. However, predictions based on historical patterns require periodic reassessment as local disease


epidemiology and healthcare utilization change (Takase & Carlin, 2012). Staffing arrangements represent one of the most important operational applications of seasonal forecasting. Hospitals may need additional pediatric nurses, physicians, respiratory therapists, or administrative personnel during anticipated periods of increased attendance. However, the appropriate allocation of staff depends on more than patient numbers. Overnight attendance may involve fewer patients but a greater proportion of severe conditions, while evening shifts may include substantial volumes of minor trauma and common infections. Research examining pediatric arrivals across different shifts supports the need for staffing decisions that incorporate patient acuity and clinical presentation (Almubarak et al., 2019). School calendars and public holidays can also influence resource planning. An investigation of pediatric emergency attendance over seven years found that Mondays generally experienced higher patient volumes, whereas certain holidays had reduced attendance. Seasonal differences were also evident, with lower attendance during summer. These patterns suggest that staffing models based exclusively on average daily volume may fail to reflect predictable changes in demand. Healthcare institutions should consider recurring calendar effects while maintaining the flexibility required for unexpected emergencies (Krinsky-Diener et al., 2017). The COVID-19 pandemic demonstrated that emergency utilization patterns can change substantially when population behavior, infection prevention practices, school attendance, and healthcare access are disrupted. Although historical seasonal models remain valuable, healthcare systems should recognize that unusual outbreaks and public health interventions can alter previously established trends. Research examining infectious disease patterns in febrile infants found changes in viral detection during pandemic-related periods, followed by renewed viral activity. Such findings emphasize the importance of incorporating current surveillance information into operational planning (Bellini et al., 2026). Healthcare equity is another important consideration. Emergency departments frequently provide care to families who face difficulties obtaining timely primary healthcare services. Socioeconomic circumstances, transportation, insurance arrangements, and geographical access may influence the decision to seek emergency treatment. Institutions should therefore interpret attendance data within the broader healthcare context rather than assuming that every low-acuity visit represents inappropriate use of emergency resources. Improving access to community healthcare may complement emergency department planning and reduce avoidable pressure on hospital services (Hasegawa et al., 2015). Operational evaluation should incorporate multiple performance indicators, including monthly attendance, peak daily volume, triage distribution, waiting times, hospital admission rates, intensive care utilization, and emergency department length of stay. Disease-specific measures can help identify whether seasonal increases arise from respiratory illness, gastrointestinal disease, trauma, or other conditions. Continuous monitoring of these indicators may support earlier recognition of emerging healthcare demands and improve the responsiveness of pediatric emergency services (Guo et al., 2023).

Future Directions and Evidence-Based Strategies for Pediatric Emergency Preparedness

Future research and healthcare planning should prioritize the development of integrated surveillance systems capable of identifying changing pediatric emergency patterns in a timely and clinically meaningful manner. Advances in electronic health records, health information systems, statistical modeling, and epidemiological monitoring provide opportunities to examine large volumes of emergency attendance data. However, reliable planning requires more than collecting total patient numbers. Future systems should evaluate specific diagnoses, age categories, severity levels, and geographical patterns to support more precise allocation of healthcare resources (Tachinardi et al., 2026). Predictive analytics represents a promising approach to improving pediatric emergency preparedness. Historical emergency attendance can be analyzed alongside seasonal indicators, school calendars, weather conditions, and infectious disease surveillance data. These variables may help identify periods when hospitals are likely to experience increased demand. Time-series research conducted in Lisbon demonstrated that multiple recurring patterns can be incorporated into forecasting models with reasonable predictive performance. Nevertheless, models should be validated using independent data and recalibrated when local healthcare conditions change (Almeida et al.,

2022). Disease-specific forecasting may provide greater clinical value than general attendance prediction alone. Different pediatric conditions have distinct seasonal distributions, and hospitals may require different resources depending on the predominant diagnoses. For example, an increase in respiratory presentations may create additional demand for oxygen therapy and respiratory monitoring, while an increase in traumatic injuries may require greater imaging and procedural capacity. Research examining condition-specific seasonal rhythms found that many pediatric presenting complaints exhibited distinct temporal patterns, supporting the development of more detailed forecasting approaches (Tachinardi et al., 2026). Future systems should incorporate age-specific analysis because infants, preschool children, school-aged children, and adolescents experience different patterns of disease. In particular, febrile infants require careful evaluation because serious bacterial infections may occur even when seasonal viral activity is high. Evidence from a recent observational study demonstrated that viral infections in young infants varied seasonally, whereas serious bacterial infections did not follow the same pattern. This supports the use of age-based clinical risk assessment alongside epidemiological information rather than allowing seasonal expectations to replace diagnostic evaluation (Bellini et al., 2026). Respiratory disease preparedness should include strengthened surveillance of asthma exacerbations and circulating respiratory pathogens. Studies conducted in Korea, Türkiye, and Italy have demonstrated that asthma-related emergency attendance varies by season and age group. These findings suggest that hospitals and community healthcare services may benefit from planning preventive asthma education and reviewing access to treatment before anticipated high-risk periods. However, preventive activities should be based on local patterns because the timing of seasonal exacerbations is not identical across countries (Won et al., 2016). Community-based preventive healthcare may reduce some potentially avoidable pediatric emergency presentations. Vaccination programs, infection prevention education, asthma management plans, access to primary healthcare, and appropriate caregiver guidance can support earlier treatment of common childhood illnesses. However, the impact of these interventions on emergency attendance should be evaluated rather than assumed. Healthcare institutions should ensure that prevention messages do not discourage families from seeking urgent assessment when children demonstrate serious symptoms or signs of deterioration (Dondi et al., 2017). Further investigation of gastrointestinal and injury-related presentations is also necessary. Seasonal changes in food exposure, water safety, recreational activities, and population movement may influence pediatric emergency demand, but these relationships are likely to differ among regions. Research examining a destination-city emergency department showed that visiting populations experienced different seasonal utilization patterns from local residents. Such findings support the inclusion of tourism, travel, and population mobility when studying emergency departments serving regions with substantial seasonal population changes (Rose et al., 2022). Healthcare organizations should also consider the integration of real-time attendance dashboards and clinical surveillance platforms. These systems could provide regular updates concerning emergency visit numbers, diagnostic categories, triage severity, waiting times, and inpatient capacity. Automated monitoring may help administrators identify unexpected increases in specific illnesses and coordinate additional resources when required. However, electronic dashboards must use accurate diagnostic coding and maintain appropriate data privacy safeguards. The value of such systems should be assessed through improvements in operational decisions and patient care (Guo et al., 2023). Artificial intelligence and machine learning may offer additional opportunities for predicting emergency demand, although their application should be supported by transparent evaluation. Predictive models may identify relationships among historical patient arrivals, environmental data, disease patterns, and hospital capacity. Nevertheless, complex algorithms can produce inaccurate forecasts when their training data do not reflect current epidemiological conditions. Future research should compare advanced predictive systems with simpler statistical approaches and evaluate whether improvements in forecasting accuracy lead to meaningful operational benefits (Takase & Carlin, 2012). Future investigations should distinguish emergency attendance, inpatient admission, and intensive care utilization. These outcomes represent different levels of healthcare demand and should be analyzed independently. A hospital may experience

increased respiratory infection attendance without a proportional rise in admissions, while a relatively small number of severe cases may substantially increase demand for intensive care. Evaluations should therefore incorporate age-specific hospitalization rates, disease severity, treatment requirements, and patient outcomes to support more complete healthcare planning (Solmaz & Orhan, 2026). Multicenter research is needed to understand geographical variation in pediatric emergency patterns. Findings from hospitals in temperate regions may not apply directly to tropical environments, rural communities, or healthcare systems with different referral arrangements. Comparative investigations should examine the influence of climate, population density, socioeconomic circumstances, school schedules, and access to primary care. Standardized data collection could improve comparisons while allowing regional healthcare organizations to develop locally appropriate preparedness strategies (Hasegawa et al., 2015). The development of effective emergency preparedness also requires collaboration among hospital administrators, pediatric clinicians, epidemiologists, public health specialists, and community healthcare providers. These professionals can contribute information concerning disease surveillance, clinical severity, staffing needs, and preventive healthcare priorities. Coordinated planning may improve the ability of pediatric emergency departments to manage seasonal fluctuations while maintaining timely treatment for children with urgent medical conditions (Gowa et al., 2019). Ultimately, future pediatric emergency planning should be evidence-based, flexible, and responsive to changing disease patterns. Seasonal forecasting can support preparedness, but it should not replace real-time surveillance or clinical assessment. Integrating demographic analysis, disease-specific monitoring, operational evaluation, and community prevention strategies may strengthen pediatric emergency care and improve the use of healthcare resources throughout the year (Tachinardi et al., 2026).



Summary

Pediatric emergency departments experience substantial variation in patient attendance, clinical presentation, and healthcare resource requirements throughout the year. These patterns are influenced by demographic characteristics, infectious disease circulation, environmental conditions, school calendars, population movement, and access to healthcare services. Published research indicates that respiratory infections contribute to increased pediatric emergency attendance during colder months in many temperate regions, while asthma exacerbations, gastrointestinal illnesses, traumatic injuries, and other conditions demonstrate distinct seasonal distributions. However, these

patterns vary across geographical locations and patient age groups. An important distinction exists between total emergency attendance and hospital admission. High patient volume does not necessarily indicate increased disease severity, making it essential to evaluate triage classifications, treatment requirements, hospitalization rates, and intensive care utilization separately. Seasonal forecasting, electronic health record analysis, and disease-specific surveillance may improve staffing, diagnostic preparedness, and resource allocation. Nevertheless, predictive models require ongoing validation because changes in infectious disease patterns and healthcare-seeking behavior can influence future attendance. Effective pediatric emergency preparedness should combine historical epidemiological evidence with real-time monitoring, preventive healthcare, multidisciplinary planning, and equitable access to appropriate treatment. Strengthening these approaches may improve the efficiency and responsiveness of pediatric emergency services while supporting safe and timely care for children throughout the year.

References:

Almeida, H. S., Sousa, M., Mascarenhas, I., Russo, A., Barrento, M., Mendes, M., Nogueira, P., & Trigo, R. (2022). The dynamics of patient visits to a public hospital pediatric emergency department: A time-series model. Pediatric Emergency Care, 38(1), e240–e245. https://doi.org/10.1097/PEC.0000000000002235

Almubarak, H., Meckler, G., & Doan, Q. (2019). Factors and outcomes associated with paediatric emergency department arrival patterns through the day. Paediatrics & Child Health, 24(5), 323–329. https://doi.org/10.1093/pch/pxy173

Altaş, U., Söğütlü, Y., Altaş, Z. M., & Özkars, M. Y. (2023). Seasonal pattern of emergency department visits by children with asthma. Eastern Mediterranean Health Journal, 29(10), 789–795. https://doi.org/10.26719/emhj.23.114

Bellini, T., Brisca, G., Mariani, M., Ferretti, M., Lacovara, A., Pezzotta, F., Vanorio, B., Pastorino, A., Serafino, F., Lasagna, J., Matarese, S., Canzoneri, F., Piccotti, E., & Moscatelli, A. (2026). Seasonal and age-specific patterns of viral and bacterial infections in febrile infants ≤90 days presenting to a paediatric emergency department. Acta Paediatrica, 115(9), 1991–1999. https://doi.org/10.1111/apa.70599

Dondi, A., Calamelli, E., Piccinno, V., Ricci, G., Corsini, I., Biagi, C., & Lanari, M. (2017). Acute asthma in the pediatric emergency department: Infections are the main triggers of exacerbations. BioMed Research International, 2017, Article 9687061. https://doi.org/10.1155/2017/9687061

Gowa, M., Habib, I., Tahir, A., Yaqoob, U., & Junejo, S. (2019). Disease spectrum and frequency of illness in pediatric emergency: A retrospective analysis from Karachi, Pakistan. Ochsner Journal, 19(4), 340–346. https://doi.org/10.31486/toj.18.0134

Guo, L. L., Guo, L. Y., Li, J., Gu, Y. W., Wang, J. Y., Cui, Y., Qian, Q., Chen, T., Jiang, R., & Zheng, S. (2023). Characteristics and admission preferences of pediatric emergency patients and their waiting time prediction using electronic medical record data: Retrospective comparative analysis. Journal of Medical Internet Research, 25, e49605. https://doi.org/10.2196/49605

Hasegawa, K., Tsugawa, Y., Cohen, A., & Camargo, C. A., Jr. (2015). Infectious disease-related emergency department visits among children in the United States. Pediatric Infectious Disease Journal, 34(7), 681–685. https://doi.org/10.1097/INF.0000000000000704

Krinsky-Diener, M., Agoritsas, K., Chao, J. H., & Sinert, R. (2017). Predicting flow in the pediatric emergency department: Are holidays lighter? Pediatric Emergency Care, 33(5), 339–343. https://doi.org/10.1097/PEC.0000000000000714

Lipsett, S. C., Monuteaux, M. C., & Fine, A. M. (2021). Seasonality of common pediatric infectious diseases. Pediatric Emergency Care, 37(2), 82–85. https://doi.org/10.1097/PEC.0000000000001496

Rose, H. S., Wolf, B. J., Pruitt, C. M., & Kane, I. D. (2022). Pediatric ED utilization in a destination city: Demographic, clinical, and temporal trends. Pediatric Emergency Care, 38(8), e1472–e1478. https://doi.org/10.1097/PEC.0000000000002792

Solmaz, M., & Orhan, O. (2026). Evaluation of demographic and clinical characteristics of pediatric emergency department visits in southeastern Turkey, 2023. Innovative Journal of Pediatrics, 36(1), e165075. https://doi.org/10.5812/ijpediatr-165075

Tachinardi, P., Witt, R. M., Wu, G., Paulose, J. K., Wong, H. R., Dribin, T. E., Mendonca, E. A., Smith, D. F., Hogenesch, J. B., & Ruben, M. D. (2026). Condition-specific seasonal rhythms in pediatric emergencies. Scientific Reports. Advance online publication. https://doi.org/10.1038/s41598-026-78609-9

Takase, M., & Carlin, J. (2012). Modelling seasonal variations in presentations at a paediatric emergency department. Hiroshima Journal of Medical Sciences, 61(3), 51–58. https://pubmed.ncbi.nlm.nih.gov/23077863/

Won, Y. K., Hwang, T. H., Roh, E. J., & Chung, E. H. (2016). Seasonal patterns of asthma in children and adolescents presenting at emergency departments in Korea. Allergy, Asthma & Immunology Research, 8(3), 223–229. https://doi.org/10.4168/aair.2016.8.3.223

Rose, H. S., Wolf, B. J., Pruitt, C. M., & Kane, I. D. (2022). Pediatric ED utilization in a destination city: Demographic, clinical, and temporal trends. Pediatric Emergency Care, 38(8), e1472–e1478. https://doi.org/10.1097/PEC.0000000000002792

Almeida, H. S., Sousa, M., Mascarenhas, I., Russo, A., Barrento, M., Mendes, M., Nogueira, P., & Trigo, R. (2022). The dynamics of patient visits to a public hospital pediatric emergency department: A time-series model. Pediatric Emergency Care, 38(1), e240–e245. https://doi.org/10.1097/PEC.0000000000002235

Guo, L. L., Guo, L. Y., Li, J., Gu, Y. W., Wang, J. Y., Cui, Y., Qian, Q., Chen, T., Jiang, R., & Zheng, S. (2023). Characteristics and admission preferences of pediatric emergency patients and their waiting time prediction using electronic medical record data: Retrospective comparative analysis. Journal of Medical Internet Research, 25, e49605. https://doi.org/10.2196/49605

Ramgopal, S., Phamduy, T. T., Keating, E. M., Shapiro, D. J., Badaki-Makun, O., Eltorki, M., Bergmann, K. R., Gonzalez, F., Geanacopoulos, A. T., Rees, C. A., Chaudhari, P. P., & Neuman, M. I. (2026). Trends in emergency department visits for children, 2016–2023. The Journal of Pediatrics, 299, 115286. https://doi.org/10.1016/j.jpeds.2026.115286

Lipsett, S. C., Monuteaux, M. C., & Fine, A. M. (2021). Seasonality of common pediatric infectious diseases. Pediatric Emergency Care, 37(2), 82–85. https://doi.org/10.1097/PEC.0000000000001496

Wrotek, A., Czajkowska, M., & Jackowska, T. (2020). Seasonality of respiratory syncytial virus hospitalization. Advances in Experimental Medicine and Biology, 1279, 93–100. https://doi.org/10.1007/5584_2020_503

Kusma, J. D., Macy, M. L., Kociolek, L. K., Davis, M. M., & Ramgopal, S. (2023). Seasonality in respiratory syncytial virus hospitalizations and immunoprophylaxis. JAMA Health Forum, 4(6), e231582. https://doi.org/10.1001/jamahealthforum.2023.1582

Gupta, P., Beam, B. W., & Rettiganti, M. (2016). Temporal trends of respiratory syncytial virus-associated hospital and ICU admissions across the United States. Pediatric Critical Care Medicine, 17(8), e343–e351. https://doi.org/10.1097/PCC.0000000000000850

Won, Y. K., Hwang, T. H., Roh, E. J., & Chung, E. H. (2016). Seasonal patterns of asthma in children and adolescents presenting at emergency departments in Korea. Allergy, Asthma & Immunology Research, 8(3), 223–229. https://doi.org/10.4168/aair.2016.8.3.223

Johnston, N. W., Johnston, S. L., Norman, G. R., Dai, J., & Sears, M. R. (2006). The September epidemic of asthma hospitalization: School children as disease vectors. Journal of Allergy and Clinical Immunology, 117(3), 557–562. https://doi.org/10.1016/j.jaci.2005.11.034

Johnston, N. W., Johnston, S. L., Duncan, J. M., Greene, J. M., Kebadze, T., Keith, P. K., Roy, M., Waserman, S., & Sears, M. R. (2005). The September epidemic of asthma exacerbations in children: A search for etiology. Journal of Allergy and Clinical Immunology, 115(1), 132–138. https://doi.org/10.1016/j.jaci.2004.09.025

Sears, M. R., & Johnston, N. W. (2007). Understanding the September asthma epidemic. Journal of Allergy and Clinical Immunology, 120(3), 526–529. https://doi.org/10.1016/j.jaci.2007.05.047

Shah, M. P., Tate, J. E., Steiner, C. A., & Parashar, U. D. (2016). Decline in emergency department visits for acute gastroenteritis among children in 10 US states after implementation of rotavirus vaccination, 2003 to 2013. The Pediatric Infectious Disease Journal, 35(7), 782–786. https://doi.org/10.1097/INF.0000000000001175

McCain, J. E., Bridgmon, A. E., King, W. D., & Monroe, K. (2023). Children's injury database: Development of an injury surveillance system in a pediatric emergency department. Injury Epidemiology, 10(Suppl. 1), Article 40. https://doi.org/10.1186/s40621-023-00443-8

Wells, J. M., Rodean, J., Cook, L., Sills, M. R., Neuman, M. I., Kornblith, A. E., Jain, S., Hirsch, A. W., Goyal, M. K., Fleegler, E. W., DeLaroche, A. M., Aronson, P. L., & Leonard, J. C. (2022). Injury-related pediatric emergency department visits in the first year of COVID-19. Pediatrics, 150(4), e2021054545. https://doi.org/10.1542/peds.2021-054545

Bernstein, A. S., Sun, S., Weinberger, K. R., Spangler, K. R., Sheffield, P. E., & Wellenius, G. A. (2022). Warm season and emergency department visits to U.S. children's hospitals. Environmental Health Perspectives, 130(1), 017001. https://doi.org/10.1289/EHP8083

Janke, A. T., Mangus, C. W., Fung, C. M., Kamdar, N., Macy, M. L., Nypaver, M. M., & Kocher, K. E. (2023). Emergency department care for children during the 2022 viral respiratory illness surge. JAMA Network Open, 6(12), e2346769. https://doi.org/10.1001/jamanetworkopen.2023.46769

Michelson, K. A., Monuteaux, M. C., Stack, A. M., & Bachur, R. G. (2012). Pediatric emergency department crowding is associated with a lower likelihood of hospital admission. Academic Emergency Medicine, 19(7), 816–820. https://doi.org/10.1111/j.1553-2712.2012.01390.x

Gross, T. K., Lane, N. E., & Timm, N. L. (2023). Crowding in the emergency department: Challenges and best practices for the care of children. Pediatrics, 151(3), e2022060972. https://doi.org/10.1542/peds.2022-060972

Foster, A. A., Eisenberg, J., Crady, R., Hewes, H. A., & Remick, K. E. (2025). Changes in pediatric inpatient capabilities and emergency department pediatric readiness. JAMA Network Open, 8(5), e258277. https://doi.org/10.1001/jamanetworkopen.2025.8277

Newgard, C. D., Rakshe, S., Salvi, A., Lin, A., Cook, J. N. B., Gausche-Hill, M., Kuppermann, N., Goldhaber-Fiebert, J. D., Burd, R. S., Malveau, S., Jenkins, P. C., Stephens, C. Q., Glass, N. E., Hewes, H., Mann, N. C., Ames, S. G., Fallat, M., Jensen, A. R., Ford, R. L., ... Remick, K. E. (2024). Changes in emergency department pediatric readiness and mortality. JAMA Network Open, 7(7), e2422107. https://doi.org/10.1001/jamanetworkopen.2024.22107

Newgard, C. D., Lin, A., Goldhaber-Fiebert, J. D., Remick, K. E., Gausche-Hill, M., Burd, R. S., Malveau, S., Cook, J. N. B., Jenkins, P. C., Ames, S. G., Mann, N. C., Glass, N. E., Hewes, H. A., Fallat, M., Salvi, A., Carr, B. G., McConnell, K. J., Stephens, C. Q., Ford, R., ... Kuppermann, N. (2024). State and national estimates of the cost of emergency department pediatric readiness and lives saved. JAMA Network Open, 7(11), e2442154. https://doi.org/10.1001/jamanetworkopen.2024.42154

Remick, K. E., Foster, A. A., Jensen, A. R., Williams, R. F., Stone, E., Joseph, M., Conners, G. P., Brown, K., & Gausche-Hill, M. (2026). Pediatric readiness in the emergency department: Policy statement. Pediatrics, 157(2), e2025075318. https://doi.org/10.1542/peds.2025-075318

Ames, S. G., Davis, B. S., Marin, J. R., Fink, E. L., Olson, L. M., Gausche-Hill, M., Kahn, J. M., & Kuppermann, N. (2019). Emergency department pediatric readiness and mortality in critically ill children. Pediatrics, 144(3), e20190568. https://doi.org/10.1542/peds.2019-0568